

Biting



Biting is a common but upsetting behavior of toddlers and two year olds. Because it is upsetting and potentially dangerous, it is important for parents to address biting when it occurs.

When a child bites another child

Intervene immediately between the child who bit and the bitten child. Stay calm; don't overreact, yell or give a lengthy explanation.


Use your voice and expression to show that biting is not acceptable. Look into the child's eyes and say calmly but firmly "No biting people." Point out how the biter's behavior affected the other person. "You hurt him and he's crying." Encourage the child who was bitten to tell the biter "You hurt me." Encourage the child who bit to help the other child by getting the ice pack, etc.

Offer the bitten child comfort and first aid. Wash broken skin with warm water and soap. Observe standard precautions if there is bleeding. Apply an ice pack or cool cloth to help prevent swelling. If the bitten child is a guest, tell the parents what happened. Suggest the bitten child be seen by a health care provider if the skin is broken or there are any signs of infection (redness or swelling).

Preventing biting

Reinforce desired behavior. Notice and acknowledge when you like what your child is doing, especially for showing empathy or social behavior, such as patting a crying child, offering to take turns with a toy or hugging gently. Do not label, humiliate or isolate a child who bites.

Discourage play which involves "pretend" biting, or seems too rough and out of control. Help the child learn to communicate by using words.



Provided by California Childcare Health Program
For more information, please contact:
Healthline 1-800-333-3212

Distributed by:

Why do children bite and what can we do?

Children bite for many different reasons, so in order to respond effectively it's best to try and find out why they are biting. Keep notes over several days on when, where and why your child bites. This may help you see a pattern and understand how to intervene.

If your child experiments by biting, immediately say "no" in a firm voice, and give him a variety of toys to touch, smell and taste and encourage sensory-motor exploration.

If your child has teething discomfort, provide cold teething toys or safe, chewy foods.



If your child is becoming independent, provide opportunities to make age-appropriate choices and have some control (the bread or the cracker, the yellow or the blue ball), and notice and give positive attention as new self-help skills and independence develop.

If your child is using muscles in new ways, provide a variety of play materials (hard/soft, rough/smooth, heavy/light) and plan for plenty of active play indoors and outdoors.

If your child is learning to play with other children, try to guide behavior if it seems rough (take the child's hand and say, "Touch Jorge gently—he likes that") and reinforce pro-social behavior (such as taking turns with toys or patting a crying child).

If your child is frustrated in expressing his/her feelings, needs and wants, state what she is trying to communicate ("you feel mad when Ari takes your truck" or "you want me to pay attention to you").

If your child is threatened by new or changing situations such as a parent returning to work, a new baby, or parents separating, provide special nurturing and be as warm and reassuring as possible, and help him talk about feelings even when he says thing like "I hate my new baby."

- Consult with a professional if your child seems to be acting out due to unusual stress.
- If the child continues biting over several weeks or does not seem to care about the consequences, seek professional help. It is unusual for a preschool age (3-5 year old) child to continue to bite and he/she needs to be evaluated for developmental concerns.

by Cheryl Oku, Infant/Toddler Specialist